

《单词百科:ischemiareperfusion是什么意思?ischemiareperfusion怎么发音? ischemiareperfusion的解释和用法》

英语单词ischemiareperfusion是什么意思?ischemiareperfusion怎么读?ischemiareperfusion怎么发音?简答网为您整理了ischemiareperfusion的解释、用法、例句、词组等相关学习资料。下面跟小编一起来看看吧!



ischemiareperfusion的意思

1、缺血再灌注

ischemiareperfusion的双语例句

1、Results: TTC in the ischemia group was larger than that in Astragalus Membranaceus group and the ischemiareperfusion group.

结果：再灌注组和黄芪治疗组的梗死面积小于单纯缺血组；

2、The results showed that the treatment with verapamil before ischemia produced significant protection against gastric injury in rats induced by ischemiareperfusion.

结果表明：戊脉安无论是胃内给药还是肌内注射，均能明显地降低胃粘膜损伤，起到保护作用。

3、Methods We duplicated the model of limb ischemiareperfusion injury in rabbit, than the contents of LDH, CD, MDA, SOD were observed, and differences of grounds were analysed.

方法复制家兔后肢缺血再灌注（IR）的损伤模型，取不同组的血浆测定其LDH、CD、MDA、SOD的含量。

4、Results: After renal ischemiareperfusion injury in rats, the level of MDA was lower and the activity of SOD was higher in puerarin group than those in the control.

结果：急性肾缺血再灌注损伤后，葛根素组大鼠MDA含量显著下降，SOD活性明显提高，肾组织病理改变明显轻于对照组；

5、Objective: To investigate the effects of exogenous epidermal growth factor (EGF) on gut function and mucosal permeability in rats after gut ischemiareperfusion (IR) insult.

目的：观察大鼠肠缺血再灌注后外源性表皮生长因子（EGF）对肠黏膜通透性和肠、肝、肾功能改变的影响。

6、Objective To establish an appropriate model of focal ischemiareperfusion in experimental chronic diabetic rats and comparatively analyze the factors influencing successful preparation of model.

目的建立合适的实验性慢性糖尿病大鼠局灶性脑缺血再灌注模型，并比较分析影响模型制备成功的因素。

7、Objective To investigate the effect of Shenfu injection (SFI) on myocardial ischemiareperfusion injury (IRI) in mitral valve replacement (MVR).

目的观察参附注射液（SFI）对二尖瓣置换术（MVR）中心肌的保护作用。

8、The possibility of using red blood cells (RBC) as the carrier of superoxide dismutase (SOD) was studied in this paper. SOD was encapsulated into rabbit erythrocytes (RBC-SOD), and was used in the study of cerebral ischemiareperfusion (I-R).

研究用红细胞（RBC）作为超氧化物歧化酶（SOD）载体的可行性，将包载SOD的RBC（RBC-SOD）用于家兔脑缺血-再灌注（I-R）模型研究。

9、Conclusion Dipyridamole can improve hepatic microcirculation and has a protective effect on hepatic ischemiareperfusion injury.

结论潘生丁预处理可明显改善肝微循环，减轻肝缺血再灌注损伤。

10、 Objective: To investigate the effect of electrical stimulation of hypothalamus paraventricular nucleus (PVN) on gastric ischemiareperfusion injury (GIRI) in rats.

目的：观察电刺激下丘脑室旁核（PVN）对大鼠胃缺血再灌注损伤（GIRI）的影响及形态学变化。

以上是简答网为您整理的ischemiareperfusion怎么读的相关信息，希望对大家有一定的帮助。查看更多关于ischemiareperfusion的用法、ischemiareperfusion的释义、ischemiareperfusion的相关详情请点击：<https://dict.jiandongshicai.cn/ischemiareperfusion>

